ALLERGY, VITAMINS

AN UPDATE OF THE BARKER HYPOTHESIS

27.02.2007

Do you know the Barker hypothesis? It says that early life events may induce later disease. A new study in the <u>Eur | Clin Nutr</u> now reports

Children whose mothers had a 25(OH)-vitamin D concentration in pregnancy >75 nmol/l had an increased risk of eczema on examination at 9 months (OR 3.26, 95% CI 1.15–9.29, P=0.025) and asthma at age 9 years (OR 5.40, 95% CI, 1.09–26.65, P=0.038) compared to children whose mothers had a concentration of <30 nmol/l.

These are quite remarkable results. I wonder only why two forthcoming studies <u>click</u> + <u>click</u> using food frequency questionnaires will describe only a reduction of respiratory infection rates (which is not too bad in the context of the hygiene hypothesis) with no effects on allergy.

Food intake measures are probably not good enough for internal dose <u>estimates</u> as main source of vitamin D production is sunlight, isn´t it?

Addendum

12 March 2007 – the long awaited <u>AJCN</u> paper is now out. I have already feared that stratification by FFQ is only partially relevant. Indeed, the authors could have used also college graduate level which increased with intake catgeories 1-2-3-4 from 53-74-80-82% with all the problems by <u>SES</u> dependency (and not corrected for in model 2+3 in table 2 or model A...D in table 3). The discussion seems to be also far away from being an objective report on the immunology of vitamin D actions...

CC-BY-NC Science Surf 27.02.2007, access 18.10.2025 ☐